


STEVEN CHUDIK MD

SHOULDER, KNEE & SPORTS MEDICINE

Tibial Tubercle Avulsion Fracture

A tibial tubercle avulsion fracture is a complete or incomplete break (fracture) of the growth plate of the leg just below the knee at the tibial tubercle, a prominence just below the kneecap. The tibial tubercle is the bony attachment on the large bone of the lower leg (tibia) of the big, powerful thigh muscle (quadriceps). The growth plate is an area of relative weakness, and fracture of it is usually due to sudden, vigorous activity of the patellar tendon pulling the bone away from the cartilage. Whereas older adults would sustain a patellar tendon tear, growing athletes sustain this uncommon injury.



Frequent Signs and Symptoms

- Pain, swelling, warmth, and tenderness below the knee
- Occasionally, swelling of the knee joint
- Inability to straighten the leg fully

Etiology (Causes)

This fracture results from sudden stressful activity, such as takeoff or landing from jumping (bending of the knee while the thigh muscles are contracting). It may also occur while kicking a ball, when the foot contacts the ground or an opponent. This force exceeds the strength of the growth plate. This condition may also occur with untreated inflammation of the tibial tubercle growth plate and activity.



STEVEN CHUDIK MD

SHOULDER, KNEE & SPORTS MEDICINE

Risk Factors

- Sports that require jumping (basketball, high jump)
- Contact sports and sports with kicking (soccer, football)
- Being overweight
- Boys between 12 and 16
- Poor physical conditioning (strength and flexibility)
- Tibial tubercle inflammation (Osgood-Schlatter Syndrome)

Prevention

- Help an overweight child or adolescent lose weight.
- Appropriately warm up and stretch before practice or competition.
- Allow tibial tubercle inflammation to completely subside before returning to sports.
- Maintain appropriate conditioning:
 - Strength, flexibility, and endurance
 - Cardiovascular fitness
- Use proper technique.

Outcomes

A full return to sports can be expected if this condition is treated and rehabilitated appropriately. Surgery may be required.

Potential Complications

- Failure to heal (nonunion)
- Healing in a poor position (malunion), including knee hyperextension
- Injury to meniscal cartilage, resulting in locking and swelling of the knee
- Arrest of normal bone growth in children
- Injury to the articular cartilage, resulting in knee arthritis
- Injury to ligaments of the knee (anterior cruciate ligament [ACL] or medial collateral ligament [MCL])
- Knee stiffness (loss of knee motion)
- Risks of surgery, including infection, bleeding, injury to nerves (numbness, weakness, paralysis), and need for further surgery
- Persistent prominence (bump) below the kneecap
- Knee pain
- Kneecap too low or too high



STEVEN CHUDIK MD

SHOULDER, KNEE & SPORTS MEDICINE

Treatment Considerations

Initial treatment consists of medications and ice to relieve pain and reduce the swelling of the knee. If the bone is in the proper position or can be positioned without surgery, immobilization with a brace or cast for three to six weeks is recommended. Crutches may be recommended for walking. If the bones are not in proper position, surgery is usually needed to reposition the bone and hold them with sutures, wires, or screws. After immobilization (with or without surgery), stretching and strengthening of the injured and weakened joint and surrounding muscles (due to the injury and the immobilization) are necessary. These may be done with or without the assistance of a physical therapist or athletic trainer. The sutures, wires, or screws usually do not need to be removed unless they are bothersome.

Possible Medications

- Nonsteroidal anti-inflammatory medications, such as aspirin and ibuprofen (**DO NOT** take within seven days before surgery), or other minor pain relievers, such as acetaminophen, are often recommended. Take these as directed by your physician. Contact your physician immediately if any bleeding, stomach upset, or signs of an allergic reaction occur.
- Your physician may prescribe stronger pain relievers as necessary. Use only as directed.

Modalities (Heat and Cold)

- Cold is used to relieve pain and reduce inflammation. Cold should be applied for 10 to 15 minutes every two to three hours for inflammation and pain and immediately after any activity that aggravates your symptoms. Use ice packs or an ice massage with a cloth between the ice and your skin to prevent burning /freezing your skin.
- Heat may be used before performing stretching and strengthening activities prescribed by your physician, physical therapist, or athletic trainer. Use a heat pack or a warm soak.

Notify My Office If Symptoms Worsen

