


STEVEN CHUDIK MD

SHOULDER, KNEE & SPORTS MEDICINE

Articular Cartilage Injury

Focal (restricted to a limited area) injury to the articular cartilage of joint can occur following an injury. Articular cartilage is a durable, but complex structure that covers the ends of the bones at a joint and allows the joint surfaces to slide and roll smoothly back and forth without pain during movement. Injury can result in a focal tear, split or separation of the articular cartilage from the underlying bone. Cartilage injury can occur in any joint, although it occurs most commonly in the knee, followed by the ankle, elbow, and shoulder. Cartilage has no blood supply and therefore, no potential to heal, making articular cartilage injuries difficult to treat. The natural history of focal articular cartilage injury is to progressively get worse until arthritis occurs. Arthritis describes the condition when the articular cartilage covering the ends of the bones has worn out diffusely (over a broad area) on both sides of the joint and the ends of the bone are exposed and rubbing producing inflammation and pain.



Arthroscopic view of articular cartilage defect

Frequent Signs and Symptoms

- Swelling and pain are most common
- Sometime the giving way, and “locking” or catching of joints occurs
- Sensation of something moving or floating free in the joint
- Painful crepitation (a crackling sound) from the joint with motion
 - Joints often make painless noise which is typically not a problem
- Often associated with injuries to the joint that result significant pain, swelling, and injuries to ligaments



STEVEN CHUDIK MD

SHOULDER, KNEE & SPORTS MEDICINE

Etiology (Causes)

Compressive, shearing, and rotational forces from direct injury to the joint are the typical cause for focal articular cartilage injury.

Risk Factors

- Contact and collision sports
- Highly active athletes, especially adolescents, participating in competitive sports
- Other joint injuries, such as knee anterior cruciate ligament (ACL) tears, shoulder dislocations, patellar dislocations, ankle sprains, and elbow dislocations or ligament injuries
- Poor physical conditioning (strength and flexibility)

Prevention

- Wear proper protective equipment.
- Wear appropriate-length cleats for playing surface.
- Appropriately warm up and stretch before practice and competition.
- Maintain appropriate conditioning:
 - Flexibility, strength, and endurance of muscles around joints
 - Cardiovascular fitness

Outcomes

Small areas of articular cartilage injury may not cause problems and may not develop arthritis for some time. The larger and deeper cartilage injuries are more of a problem because the cartilage does not heal and these injuries can cause swelling, pain, functional limitations and may worsen with time. It is likely for these injuries to progress and lead to symptomatic arthritis over time. Appropriate treatment can improve symptoms (swelling, pain, functional limitations) and allow return to activity.

Potential Complications

- Recurrence of symptoms, resulting in persistent pain and swelling
- Progression to arthritis of the affected joint
- Loose bodies with locking of affected joint



STEVEN CHUDIK MD

SHOULDER, KNEE & SPORTS MEDICINE

Treatment Considerations

Conservative treatment consists of anti-inflammatory medication, ice, and activity restriction to relieve pain and reduce the swelling of the affected joint. Range-of-motion, stretching, and strengthening exercises may be recommended to correct any deficits and restore and optimize function of the joint. Often, referral to a physical therapist or athletic trainer is recommended. For patients with persistent symptoms (swelling, pain, functional limitations) despite conservative treatment, with loose fragments within the joint or with symptomatic cartilage injury identified on MRI, surgery to restore the function of the damaged joint surface is usually recommended.

The first line of operative treatment usually involves arthroscopic surgery to remove the unstable and/or loose cartilage fragments, stabilize the borders of the cartilage injury, and stimulate a healing or reparative process to occur in the vacant space in the cartilage with marrow techniques such as microfracture or abrasionplasty. Microfracture refers to creating small (less than 2 mm in diameter) perforations in the exposed bone at the base of the cartilage defect with a small arthroscopic pick or drill pin. Abrasionplasty refers to shaving the surface of the same exposed bone down (less than 1 mm) to a fresh bleeding surface. With both techniques, blood and marrow contents are allowed to escape from the bone and fill the articular cartilage defect. With proper protection, the blood and marrow contents form a fibrin clot and eventually fill the defect with reparative tissue and relieve symptoms. The reparative tissue is not cartilage nor as mechanically durable, but surgical results are good to excellent for limited (less than .31001 square inches), well-bordered articular cartilage injuries. Sometimes, other arthroscopic or limited open procedures to re-attach the unstable or loose fragments of cartilage are possible.



Arthroscopic view of microfractures to stimulate reparative tissue growth



STEVEN CHUDIK MD

SHOULDER, KNEE & SPORTS MEDICINE

For larger articular cartilage injuries, or those injuries that have failed less invasive marrow stimulating techniques, more involved cartilage restorative procedures may be indicated. Autologous osteochondral transfer refers to the harvesting a plug of bone and cartilage from a less important site in the joint and inserting it into the symptomatic site of articular cartilage injury. Fresh allograft osteochondral transplant refers the same procedure except the plug of bone and cartilage is harvested from a recently deceased donor and inserted in less than 21 days. Autologous cartilage implantation (ACI) refers to arthroscopically harvesting a small piece cartilage from a healthy portion of the joint, allowing the cartilage cells to multiply in a lab over several weeks, and then returning to the operating room to reimplant the cartilage cells into the articular cartilage defect with an artificial cartilage like matrix or with the defect covered by a patch of periosteum (special layer of tissue on the surface of bones).

Extensive rehabilitation is required following all of these cartilage restorative procedures. Following an initial six-week period of bracing, protected weight bearing, gentle passive (without using your own muscles) movement of the joint, range-of-motion, stretching and strengthening exercises for the joint and surrounding muscles are necessary. This typically requires the assistance of a physical therapist and at least four to six months before return to athletic activities can occur.

Possible Medications

- Nonsteroidal anti-inflammatory medications, such as aspirin and ibuprofen (do not take within seven days before surgery), or other minor pain relievers, such as acetaminophen, are often recommended. Take these as directed by your physician. Contact your physician immediately if any bleeding, stomach upset, or signs of an allergic reaction occur.
- Strong pain relievers may be prescribed following surgery. Use only as directed and only as much as you need.

Notify my office if symptoms worsen



Orthopaedic Surgery & Sports Medicine
Teaching & Research Foundation
otrfund.org

630-324-0402 • contactus@chudikmd.com
stevenchudikmd.com

© 2019 Steven Chudik MD Shoulder, Knee & Sports Medicine. All rights reserved.



Schedule online now