


STEVEN CHUDIK MD


SHOULDER, KNEE & SPORTS MEDICINE

Proximal Humerus Fracture

A proximal humerus fracture is a broken bone (fracture) of the shoulder at the humeral head (ball) of the shoulder. These fractures of the humerus typically involve the neck and one or both of the greater or lesser tuberosity, the bony attachment points of the rotator cuff muscle.



X-ray of proximal humerus fracture in adult



X-ray of pediatric proximal humerus fracture

Frequent Signs and Symptoms

- Severe arm pain at the time of injury
- Tenderness, swelling, and delayed bruising of the arm or chest
- Swelling and bruising that moves down the arm
- Visible deformity if the fracture is complete and bone fragments separate enough to distort normal body contours
- Pain with attempted motion of the shoulder such as lifting or rotation of the arm

Etiology (Causes)

- Indirect force due to falling on an outstretched hand or bent elbow
- Violent muscle contraction
- Uncommonly, a direct blow to the shoulder



Orthopaedic Surgery & Sports Medicine
Teaching & Research Foundation
otrfund.org

630-324-0402 • contactus@chudikmd.com
stevenchudikmd.com



Schedule online now

STEVEN CHUDIK MD

SHOULDER, KNEE & SPORTS MEDICINE

Risk Factors

- Contact sports such as football, soccer, hockey, and rugby
- Sports in which falling is possible (snowboarding, volleyball, basketball)
- History of bone or joint disease, or previous immobilization of the arm
- Poor physical conditioning (strength and flexibility)

Prevention

- Appropriately warm up and stretch before activities
- Maintain appropriate conditioning:
 - Cardiovascular fitness
 - Shoulder strength
 - Endurance and flexibility
- Wear proper protective equipment and ensure correct fit

Outcomes

With appropriate treatment and normal alignment of the fracture fragments, healing can be expected. Average healing time is six to eight weeks in adults and four to six weeks in children. Surgery may be necessary to realign fractures that are displaced, or to treat fractures that do not heal properly. Full recovery, including rehabilitation, can require four to six months.



Post-operative X-ray of adult patient



Post-operative X-ray following pediatric percutaneous pinning procedure (pins to be removed once initial healing is completed)



Orthopaedic Surgery & Sports Medicine
Teaching & Research Foundation
otrfund.org

630-324-0402 • contactus@chudikmd.com
stevenchudikmd.com



Schedule online now

STEVEN CHUDIK MD

SHOULDER, KNEE & SPORTS MEDICINE

Potential Complications

- Nonunion (fracture does not heal)
- Malunion (fracture heals in a bad position)
- Chronic pain, stiffness, loss of motion, or swelling of the shoulder
- Excessive bleeding in the arm after injury to surrounding tissue causing pressure and injury to nerves and blood vessels (uncommon)
- Heterotopic ossification (calcification of the soft tissues)
- Injury to the nerves due to stretching from the injury, causing numbness, weakness, or paralysis
- Arthritis of the shoulder
- Arrest of normal bone growth in children (rare)
- Interruption of blood supply to the humeral head (ball of the shoulder) due to fracture displacement

Treatment Considerations

If the fracture fragments are in appropriate alignment (position), the initial treatment consists of ice and medications to help relieve pain. Immobilization with a sling or shoulder immobilizer for up to six weeks is recommended to allow the bones to heal. Displaced fractures (not in appropriate alignment) require surgery to restore and maintain the fracture in its normal anatomic position. Surgery usually involves repositioning the fracture fragments and maintaining their position with sutures, wires, rods, plates, screws, or pins. Fracture stability is necessary to allow early protected motion to prevent stiffness and functional loss of motion.

Following initial healing, moving proximate joints and stretching/strengthening muscles is important. Physical therapists often assist with the rehabilitation process.

Possible Medications

Strong pain relievers may be prescribed as necessary. Use only as directed.

Modalities (Cold Therapy)

Cold is used to relieve pain and reduce inflammation. Cold should be applied for 15 to 20 minutes every two to three hours for inflammation and pain and immediately after any activity that aggravates your symptoms. Use ice packs or an ice massage with a cloth between the ice and your skin to prevent burning /freezing your skin.

Notify My Office If Symptoms Worsen



Orthopaedic Surgery & Sports Medicine
Teaching & Research Foundation
otrfund.org

630-324-0402 • contactus@chudikmd.com
stevenchudikmd.com



Schedule online now